

WHAT IS CLAIMED IS:

1. An isolated or recombinant polynucleotide encoding an antigenic polypeptide comprising at least 17 contiguous amino acids from the mature polypeptide from SEQ ID NO: 2 or 4.
5
2. The polynucleotide of Claim 1, encoding a mature polypeptide from SEQ ID NO: 2 or 4.
- 10 3. The polynucleotide of Claim 1, which hybridizes at 55° C, less than 500 mM salt, and 50% formamide to the coding portions of SEQ ID NO: 1 or 3.
- 15 4. The polynucleotide of Claim 3, comprising at least 35 contiguous nucleotides of the coding portion of SEQ ID NO: 1 or 3.
5. An expression vector comprising the polynucleotide of Claim 1.
6. A host cell containing the expression vector of Claim 5,
20 including a eukaryotic cell.
7. A method of making an antigenic polypeptide comprising expressing a recombinant polynucleotide of Claim 1.
- 25 8. A method for forming a duplex with a polynucleotide of Claim 1,
portion of SEQ ID NO: 1 or 3; thereby forming said duplex.

9. A kit for the detection of a polynucleotide of Claim 1, comprising a polynucleotide that hybridizes, under stringent hybridization conditions, to at least 17 contiguous nucleotides of a polynucleotide of Claim 1.

5 10. The kit of claim 9, wherein said probe is detectably labeled.

11. A binding compound comprising an antibody binding site which specifically binds to at least 17 contiguous amino acids from SEQ ID NO: 2 or 4.

10

12. The binding compound of Claim 11, wherein:

a) said antibody binding site is:

1) specifically immunoreactive with a polypeptide of SEQ ID NO: 2 or 4;

15 2) raised against a purified or recombinantly produced human IL-B50 protein; or

3) in a monoclonal antibody, Fab, or F(ab)2; or

b) said binding compound is:

1) an antibody molecule;

20 2) a polyclonal antiserum;

3) detectably labeled;

4) sterile; or

5) in a buffered composition.

25 13. A method using the binding compound of Claim 11, comprising

14. A method of detecting IL-B50 in a biological sample, comprising:

15. forming a binding compound antigen complex.

16. forming a binding compound antibody sample, wherein said antibody sample

from a human, and wherein said binding compound is an antibody

15. A detection kit comprising said binding compound of Claim 12,
and:

5 a) instructional material for the use of said binding compound for said
detection; or

b) a compartment providing segregation of said binding compound.

16. A substantially pure or isolated antigenic polypeptide, which
binds to said binding composition of Claim 11, and further comprises at
10 least 17 contiguous amino acids from SEQ ID NO: 2 or 4.

17. The polypeptide of Claim 16, which:

15 a) comprises at least a fragment of at least 25 contiguous amino acid
residues from a primate IL-B50 protein;

b) is a soluble polypeptide;

c) is detectably labeled;

d) is in a sterile composition;

e) is in a buffered composition;

f) binds to a cell surface receptor;

20 g) is recombinantly produced; or

h) has a naturally occurring polypeptide sequence.

18. The polypeptide of Claim 17, which comprises at least 17
contiguous amino acids of SEQ ID NO: 2 or 4.

25

SEARCHED _____ SERIALIZED _____ INDEXED _____

antagonist of a primate IL-B50.

20. The method of Claim 19, wherein:

- a) said contacting is in combination with an agonist or antagonist of IL-7; or
- b) said contacting is with an antagonist, including binding composition comprising an antibody binding site which specifically binds an IL-B50.